=> fil reg

FILE 'REGISTRY' ENTERED AT 09:37:56 ON 19 NOV 2007 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT. PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ${\tt ZIC/VINITI}$ data file provided by InfoChem.

STRUCTURE FILE UPDATES: 18 NOV 2007 HIGHEST RN 954747-20-7 DICTIONARY FILE UPDATES: 18 NOV 2007 HIGHEST RN 954747-20-7

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 29, 2007

Please note that search-term pricing does apply when conducting ${\tt SmartSELECT}$ searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

http://www.cas.org/support/stngen/stndoc/properties.html

NODE ATTRIBUTES:
CONNECT IS E1 RC AT 5
DEFAULT MLEVEL IS ATOM
GGCAT IS SAT AT 5
GGCAT IS SAT AT 7
GGCAT IS SAT AT 7
GGCAT IS SAT AT 19
DEFAULT ECLEVEL IS LIMITED

REP G1=(2-5) C VAR G2=9/10

GRAPH ATTRIBUTES: RING(S) ARE ISOLATED OR EMBEDDED NUMBER OF NODES IS 11

STEREO ATTRIBUTES: NONE
L5 21 SEA FILE=REGISTRY SSS FUL L3

100.0% PROCESSED 603 ITERATIONS SEARCH TIME: 00.00.01

21 ANSWERS

```
(FILE 'HOME' ENTERED AT 09:19:31 ON 19 NOV 2007)
     FILE 'HCAPLUS' ENTERED AT 09:20:04 ON 19 NOV 2007
               E US20070055036/PN
L1
              1 S E3
                SEL RN
     FILE 'REGISTRY' ENTERED AT 09:20:37 ON 19 NOV 2007
L2
     FILE 'LREGISTRY' ENTERED AT 09:28:55 ON 19 NOV 2007
L3
                STR
     FILE 'REGISTRY' ENTERED AT 09:34:54 ON 19 NOV 2007
              1 S L3
L4
1.5
             21 S L3 FUL
L6
              2 S L2 AND L5
                SAV L5 LOE948/A
    FILE 'HCAPLUS' ENTERED AT 09:36:16 ON 19 NOV 2007
L7
             11 S L5
     FILE 'CAOLD' ENTERED AT 09:36:50 ON 19 NOV 2007
T.R
             0 S L5
=> fil hcap
```

FILE 'HCAPLUS' ENTERED AT 09:38:02 ON 19 NOV 2007 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT. PLEASE SEE "HELP USAGETERMS" FOR DETAILS. COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 19 Nov 2007 VOL 147 ISS 22 FILE LAST UPDATED: 18 Nov 2007 (20071118/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d 17 ibib abs hitstr hitind 1-11

L7 ANSWER 1 OF 11 HCAPLUS COPYRIGHT 2007 ACS on STN ACCESSION NUMBER: 2005:1250858 HCAPLUS Full-text DOCUMENT NUMBER: 144:129033 TITLE: Alkyne Hydrosilylation Catalyzed by a Cationic

Ruthenium Complex: Efficient and General Trans

10577,948 3

Addition

AUTHOR(S): Trost, Barry M.; Ball, Zachary T.

CORPORATE SOURCE: Department of Chemistry, Stanford University,

Stanford, CA, 94305-5080, USA
SOURCE: Journal of the American Chemical Society (2005),

127(50), 17644-17655

CODEN: JACSAT; ISSN: 0002-7863

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal LANGUAGE: English

OTHER SOURCE(S): CASREACT 144:129033

Hydrosilylation of 1-alkynes catalyzed by ruthenium cationic acetonitrile cyclopentadienyl half-sandwich complexes afforded α -substituted vinylsilanes; propargyl alcs. were converted to silylated allyl alcs. and β -hydroxyketones. The complex (Formule(MeCN)3)FP6 is shown to catalyze the hydrosilylation of a wide range of alkynes. Terminal alkynes afford access to α -vinylsilane products with good regioselectivity. Deuterium labeling studies indicate a clean trans addition process is at work. The same complex is active in internal alkyne hydrosilylation, where absolute selectivity for the trans addition process is maintained. Several internal alkyne substrate classes, including propargylic alcs. and α - β -alkynyl carbonyl compdes, allow regioselective vinylsilane formation. The tolerance of a wide range of silanes is noteworthy, including alkyl-, aryl-, alkoxy-, and halosilanes. This advantage is demonstrated in the direct synthesis of triene substrates for silion-tethered intramol. Diels-Alder cycloaddns.

IT 392700-16-2P

AR

RL: SPN (Synthetic preparation); PREP (Preparation) (regioselective hydrosilylation of alkynes, acetylenic alcs. and carbonyl compds. catalyzed by ruthenium cationic half-sandwich complexes)

RN 392700-16-2 HCAPLUS

CN 1-Oxa-2-silacyclopentane, 2,2-diethoxy-5-ethyl-3-ethylidene- (CA INDEX NAME)

CC 29-6 (Organometallic and Organometalloidal Compounds) Section cross-reference(s): 23

IT 108191-41-9P 193006-92-7P 219736-38-6P, 3,10-Tridecadiyne-1,13diol 392700-04-8P 392700-08-2P 392700-10-6P 392700-12-8P

392700-35-5P 447442-15-1P 489469-17-2P 556776-84-2P 556776-85-3P 556776-86-4P 556776-95-5P 609307-02-0P 609307-03-1P 609307-04-2P 609307-07-5P

609307-02-0P 609307-03-1P 609307-04-2P 609307-07-5P 797762-65-1P 797762-62-0P 797762-63-1P 797762-65-3P 797762-66-4P 797762-69-7P 797762-71-1P 869206-60-0P

797762-66-4P 797762-69-7P 797762-71-1P 869206-60-0P 873450-70-5P 873450-73-8P 873450-81-8P 873450-82-9P 873450-84-1P 873450-86-3P 873450-91-0P

873450-92-1P 873450-93-2P 873562-43-7P

RL: SPN (Synthetic preparation); PREP (Preparation) (regioselective hydrosilylation of alkynes, acetylenic alcs. and

carbonyl compds. catalyzed by ruthenium cationic half-sandwich

complexes)

REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 2 OF 11 HCAPLUS COPYRIGHT 2007 ACS on STN ACCESSION NUMBER: 2005:429420 HCAPLUS Full-text

DOCUMENT NUMBER: 142:464454

TITLE: Preparation of oxa-silacyclopentane type cyclic silicone compound in single step reaction for

organic silicone resin having alcoholic hydroxyl group with good long-term stability

Komuro, Katsuhiko; Suzuki, Hiroshi INVENTOR(S):

PATENT ASSIGNEE(S): Toagosei Co., Ltd., Japan PCT Int. Appl., 34 pp.

SOURCE: CODEN: PIXXD2 DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1 PATENT INFORMATION:

P2	PATENT NO.				KIND		DATE		APPLICATION NO.						DATE	
		2005044828					20050519		WO 2004-JP16445							00411
U:		CH, GB, KR, MX, SE, VC, VC, AM, DE,	CN, GD, KZ, MZ, SG, VN, GH, AZ, DK, PT, GW,	CO, GE, LC, NA, SK, YU, GM, BY, EE, RO, ML,	CR, GH, LK, NI, SL, ZA, KE, KG, ES, SE, MR,	CU, GM, LR, NO, SY, ZM, LS, KZ, FI, SI, NE,	MW, MD, FR, SK, SN,	DE, HU, LT, OM, TM, MZ, RU, GB, TR, TD,	DK, ID, LU, PG, TN, NA, TJ, GR, BF, TG	DM, IL, LV, PH, TR, SD, TM, HU, BJ,	DZ, IN, MA, PL, TT, SL, AT, IE, CF,	EC, IS, MD, PT, TZ, SZ, BE, IS, CG,	EE, JP, MG, RO, UA, TZ, BG, IT, CI,	EG, KE, MK, RU, UG, CH, LU,	BZ, ES, KG, MN, SC, US, ZM, CY, MC,	FI, KP, MW, SD, UZ, ZW, CZ, NL,
PRIORI:	PRIORITY APPLN. INFO.:						JP 2003-377752 JP 2004-39063					A 2	00609 1 00311 7			
											004-				1 W 2	00402

GΙ

OTHER SOURCE(S): MARPAT 142:464454

10577,948 5

$$\begin{array}{c} \text{OR}^2 \\ \text{Si-O} \\ \text{Z'-Me} \end{array}$$

AB A cyclic organic silicone compound I is produced by reacting an olefin ZRC(OH)CH3 and an alkoxysilane RISH(OR2)2 in the presence of a transition metal catalyst and an organic silicone resin having an alc. hydroxyl group is obtained by hydrolyzing and condensing the organic silicone compound I or a mixture of the organic silicone compound and a multifunctional alkoxysilane, wherein Z = C2-5 terminal alkenyl; R = Me or H; Rl = Cl-3 alkyl or alkoxyl group; R2 = Cl-3 alkyl group; and Z' = C2-5 alkylene group. Thus, 348 mmol 3-hydroxy-3-methyl-1- butene and 383 mmol triethoxysilane were reacted in the presence of divinyldisiloxane platinum complex to give a cyclic silicone compound with yield 82% and b.p. 117-120° at 13,300 Pa, 240 mmol of which was condensed with 441 mmol methyltriethoxysilane and 120 mmol hexamethyldisiloxane to give an alc. hydroxy-containing silicone resin, showing good long term stability.

IT 851667-88-4P

RL: IMF (Industrial manufacture); RCT (Reactant); PREP (Preparation); RACT (Reactant or reagent)

(preparation of cyclic organic silicone compds. for organic silicone resin having alc. hydroxyl group with good long-term stability)

RN 851667-88-4 HCAPLUS

CN 1-Oxa-2-silacyclopentane, 2,2-diethoxy-5,5-dimethyl- (CA INDEX NAME)

IT 851667-89-50P, trimethylsilyl-terminated

RL: IMF (Industrial manufacture); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)

(preparation of cyclic organic silicone compds. for organic silicone resin having alc. hydroxyl group with good long-term stability)

RN 851667-89-5 HCAPLUS

CN 1-Oxa-2-silacyclopentane, 2,2-diethoxy-5,5-dimethyl-, polymer with triethoxymethylsilane (9CI) (CA INDEX NAME)

CM

CRN 851667-88-4 CMF C9 H20 O3 Si

```
CRN 2031-67-6
     CMF C7 H18 O3 Si
      OEt
 Eto— Ši—Me
      bet
     ICM C07F007-18
     ICS C08G077-14
     37-3 (Plastics Manufacture and Processing)
     851667-88-4P
     RL: IMF (Industrial manufacture); RCT (Reactant); PREP
     (Preparation); RACT (Reactant or reagent)
        (preparation of cyclic organic silicone compds. for organic silicone resin
        having alc. hydroxyl group with good long-term stability)
     107-46-0DP, Hexamethyldisiloxane, reaction products with
     polysiloxane-silsesquioxanes 851667-89-5DP,
     trimethylsilvl-terminated
     RL: IMF (Industrial manufacture); TEM (Technical or engineered
     material use); PREP (Preparation); USES (Uses)
        (preparation of cyclic organic silicone compds. for organic silicone resin
       having alc. hydroxyl group with good long-term stability)
REFERENCE COUNT:
                               THERE ARE 4 CITED REFERENCES AVAILABLE FOR
                         4
                               THIS RECORD. ALL CITATIONS AVAILABLE IN
                               THE RE FORMAT
L7 ANSWER 3 OF 11 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER:
                         2001:847179 HCAPLUS Full-text
DOCUMENT NUMBER:
                         136:134826
TITLE:
                        Markovnikov Alkyne Hydrosilylation Catalyzed by
                        Ruthenium Complexes
AUTHOR(S):
                        Trost, Barry M.; Ball, Zachary T.
CORPORATE SOURCE:
                        Department of Chemistry, Stanford University,
                        Stanford, CA, 94305-5080, USA
SOURCE:
                        Journal of the American Chemical Society (2001),
                         123(50), 12726-12727
                        CODEN: JACSAT; ISSN: 0002-7863
PUBLISHER:
                        American Chemical Society
DOCUMENT TYPE:
                        Journal
LANGUAGE:
                        English
OTHER SOURCE(S):
                        CASREACT 136:134826
     The first general nondirected terminal alkyne hydrometalation to afford 1.1-
     disubstituted vinylsilanes with high regioselectivity is presented. Thus, 5-
     acetyloxy-1-pentyne is hydrosilylated with (OEt)3SiH in the presence of
     [Cp*Ru(MeCN)3]+[PF6]- to give 86% 5-acetyloxy-2-(triethoxysily1)-1-pentene,
     (I). The catalyst system extends to internal alkynes to give a single olefin
     geometry from an unusual trans addition The mild conditions and functional
     group tolerance- and the inclusion of alkoxysilanes-open a host of potential
     subsequent transformations toward the construction of complex target mols.
     For example, I readily couples with 3-iodoanisole in the presence of
     [(ally1)PdC1]2 to give 73% 5-acetyloxy-2-(3-methoxybenzene)-1-pentene.
    392700-16-2P 392700-32-2P
```

CM 2

ΤТ

10577.948 7

RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)

RN 392700-16-2 HCAPLUS

CN 1-Oxa-2-silacyclopentane, 2,2-diethoxy-5-ethyl-3-ethylidene- (CA INDEX NAME)

RN 392700-32-2 HCAPLUS

CN 1-Oxa-2-silacyclohex-3-ene, 2,2-diethoxy-6-ethyl-3-methyl- (CA INDEX NAME)

29-6 (Organometallic and Organometalloidal Compounds) 392700-04-8P 392700-06-0P 392700-10-6P 392700-12-8P 392700-14-0P, 8-(Triethylsilyl)-8-nonenoic acid 392700-16-2P 392700-17-3P 392700-18-4P 392700-19-5P 392700-20-8P, 2-(Triethoxysilv1)-1-tetradecen-3-o1 392700-22-0P, (Z)-Ethv1 3-(triethoxysily1)-2-pentenoate 392700-23-1P 392700-24-2P, 10-(Triethoxysily1)-10-undecen-3-yn-1-ol 392700-25-3P, (Z)-2-(Triethoxysily1)-2-tetradecene 392700-26-4P, (Z)-3-(Triethoxysilv1)-2-tetradecene 392700-27-5P 392700-28-6P 392700-29-7P 392700-30-0P 392700-31-1P 392700-32-2P 392700-33-3P 392700-34-4P 392700-35-5P, 1-Bromo-2-[3-(triethoxysilv1)-3-butenv1]benzene 392700-36-6P RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of) 34

REFERENCE COUNT: 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 4 OF 11 HCAPLUS COPYRIGHT 2007 ACS on STN ACCESSION NUMBER: 2000:422005 HCAPLUS Full-text

DOCUMENT NUMBER: 133:164180

TITLE: Functionalization of alkylalkoxysilanes. Studies

AUTHOR(S): towards annulations of diterpenoids
Harris, Paul W. R.; Woodgate, Paul D.
CORPORATE SOURCE: Department of Chemistry, The University of

Auckland, Auckland, N. Z.
SOURCE: Tetrahedron (2000), 56(24), 4001-4015

CODEN: TETRAB; ISSN: 0040-4020

PUBLISHER: Elsevier Science Ltd.

DOCUMENT TYPE: Journal LANGUAGE: English

8

OTHER SOURCE(S): CASREACT 133:164180

Ortho-(2-triethoxysilylethyl) derivs. of aryl ketones undergo oxidative desilvlation with H2O2. Tetralone derivs, have served as model substrates for 14-2-(triethoxysilylethyl)-7-oxopodocarpanes, which have been converted into 2-arylethanols in a four-step sequence (BH3.BMS, H202, Et3SiH/CF3COOH, TBAF) without isolation of intermediate products (oxasilepin, diol, trifluoroacetate). Use of the 14-(2-diethoxysilylmethyl) analog improved the overall yield significantly (75 vs. 40%). Re-oxidation of ring B with CAN gave the $\Delta 5-7$ -oxo derivative, which was converted into an enone-aldehyde by treatment with Collins' reagent. The conjugated alkene was reduced stereoselectively with SmI2, which, however, did not promote pinacol coupling of the resulting 1,5-keto aldehyde. ΙT

287919-15-7P

RL: SPN (Synthetic preparation); PREP (Preparation) (annulations of arenes via functionalization of alkylalkoxysilanes)

RN 287919-15-7 HCAPLUS

CN 2,3-Benzoxasilepin, 3,3-diethoxy-1,3,4,5-tetrahydro-1-methyl- (CA INDEX NAME)

287919-41-9P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(annulations of diterpenoids via functionalization of

alkvlalkoxvsilanes)

RN 287919-41-9 HCAPLUS

Phenanthro[1,2-e][1,2]oxasilepin-8-carboxylic acid, 13-[[(1,1-dimethylethyl)dimethylsilyl]oxy]-3,3-diethoxy-1,3,4,5,6,7,7a,8,9,10,11,11a-dodecahydro-1,8,11a-trimethyl-, methyl ester, (7aR, 8S, 11aS) - (CA INDEX NAME)

Absolute stereochemistry.

CC 30-20 (Terpenes and Terpenoids)

ΙT 287919-11-3P 287919-12-4P 287919-13-5P 287919-14-6P

287919-15-7P 287919-22-6P 287919-24-8P 287919-25-9P RL: SPN (Synthetic preparation); PREP (Preparation)

(annulations of arenes via functionalization of alkylalkoxysilanes)

142452-78-6P 287919-17-9P 287919-21-5P 287919-26-0P 287919-30-6P 287919-31-7P 287919-35-1P 287919-38-4P

287919-41-9P 287919-44-2P 287919-45-3P 287919-46-4P RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(annulations of diterpenoids via functionalization of

alkylalkoxysilanes)

REFERENCE COUNT: 39

THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD, ALL CITATIONS AVAILABLE IN THE RE FORMAT

ANSWER 5 OF 11 HCAPLUS COPYRIGHT 2007 ACS on STN ACCESSION NUMBER: 1998:176869 HCAPLUS Full-text

DOCUMENT NUMBER: 128:257472

TITLE: Cycloaddition of thermolytically generated

methoxymethylsilylene to α, β unsaturated ketones and imines

AUTHOR(S): Gehrhus, Barbara; Heinicke, Joachim; Meinel,

Susanne

Institut fur Anorganische Chemie, CORPORATE SOURCE:

Ernst-Moritz-Arndt-Universitat Greifswald, Greifswald, D-17487, Germany

Main Group Metal Chemistry (1998), 21(2), 99-104

CODEN: MGMCE8; ISSN: 0792-1241

PUBLISHER: Freund Publishing House Ltd.

DOCUMENT TYPE: Journal LANGUAGE: English

SOURCE:

Co-thermolyzes of 1,2-dimethyl-1,1,2,2-tetramethoxydisilane (1) with 1,4diheterodienes proceed with unsym. decay of 1 and formal [1+4]-cycloaddn. of methoxymethylsilylene to give, e.g., a 1-methoxy-1-methyl-1,3,2dioxasilacyclopentene or a 1,3,2-diazasilacyclopentene. Co-thermolytic conversions of 1 with unsatd. ketones or imines, however, afford mixts. of double bond isomers, 1-oxa- or 1-aza-2-silacyclopent-4-enes and 1-oxa- or 1aza-2-silacyclopent-3-enes, resp. The 4-ene isomers are the major products in case of oxa- and N-alkyl-azasilacyclopentenes, the 3-ene isomers dominate in a N-phenyl-azasilacyclopentene. The results can be explained by stepwise cycloaddns, of intermediate methoxymethylsilylene via transient oxa- or

158172-38-4P 205241-04-9P 205241-05-0P тт

RL: SPN (Synthetic preparation); PREP (Preparation)

(cycloaddn. of thermolytically generated methoxymethylsilylene to

α, β-unsatd, ketones and imines)

RN 158172-38-4 HCAPLUS

azasiliranes.

1-0xa-2-silacyclopent-4-ene, 2-methoxy-2,5-dimethyl-3-phenyl- (CA INDEX NAME)

CN

- 205241-04-9 HCAPLUS RN
- CN 1-Oxa-2-silacyclopent-4-ene, 5-ethyl-2-methoxy-2-methyl-3-phenyl-(CA INDEX NAME)

205241-05-0 HCAPLUS

CN 1-0xa-2-silacyclopent-4-ene, 2-methoxy-2-methyl-5-(1-methylethyl)-3phenvl- (CA INDEX NAME)

29-6 (Organometallic and Organometalloidal Compounds)

158172-38-4P 158172-39-5P 205241-02-7P 205241-03-8P

205241-04-9P 205241-05-0P 205241-06-1P

205241-08-3P 205241-07-2P 205241-09-4P 205241-10-7P

RL: SPN (Synthetic preparation); PREP (Preparation) (cycloaddn. of thermolytically generated methoxymethylsilylene to

 α , β -unsatd, ketones and imines)

REFERENCE COUNT: THERE ARE 26 CITED REFERENCES AVAILABLE 26 FOR THIS RECORD. ALL CITATIONS AVAILABLE

IN THE RE FORMAT

ANSWER 6 OF 11 HCAPLUS COPYRIGHT 2007 ACS on STN 1996:81130 HCAPLUS Full-text ACCESSION NUMBER:

DOCUMENT NUMBER: 124:261125

TITLE: Methoxymethylsilylene: (1+6)- and

(1+4)-cycloadditions to heterotrienes

AUTHOR(S): Heinicke, Joachim W.; Gehrhus, Barbara CORPORATE SOURCE: Institute Inorganic Chemistry, University Greifswald, Greifswald, D-17487, Germany SOURCE:

Heteroatom Chemistry (1995), 6(5), 461-8

CODEN: HETCE8; ISSN: 1042-7163

PUBLISHER: Wilev DOCUMENT TYPE: Journal LANGUAGE: English

GI

- AB Co-pyrolytic gas phase reactions of 1,2-dimethyl-1,1,2,2- tetramethoxydisilane with conjugated 1-oxatrienes in a flow-reactor furnish, via intermediate methoxymethylsilylene, mixts. of diastereoisomers of 3-phenyl-1-oxa-2-silacyclohepta-4,6-dienes, e.g., I, and 3-styryl-1-oxa-2-silacyclopent-4-enes, e.g., II, usually including smaller amts. of the corresponding 1-oxa-2-silacyclopent-3-ene isomers. In co-pyrolysis with an analogous N-isopropylazatriene, five-membered rings, e.g., III, were formed preferentially, and seven-membered isomers could not be detected by NMR spectroscopy.
- IT 174870-68-96 174870-70-38 174870-71-49
 174870-72-59 174870-74-72 174870-75-89
 174956-09-39 174956-11-79
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)
- RN 174870-68-9 HCAPLUS
- CN 1-Oxa-2-silacyclopent-4-ene, 2-methoxy-2,4,5-trimethy1-3-(2-phenyletheny1)-, [2 α ,3 α (E)]- (9CI) (CA INDEX NAME)

Relative stereochemistry. Double bond geometry as shown.

RN 174870-70-3 HCAPLUS

CN 1-Oxa-2-silacyclohepta-4,6-diene, 2-methoxy-2,6,7-trimethyl-3-phenyl-, cis- (9CI) (CA INDEX NAME)

Relative stereochemistry.

- RN 174870-71-4 HCAPLUS
- CN 1-0xa-2-silacyclohepta-4,6-diene, 2-methoxy-2,6,7-trimethyl-3-phenyl-, trans- (9CI) (CA INDEX NAME)

Relative stereochemistry.

- RN 174870-72-5 HCAPLUS
- CN 1-0xa-2-silacyclopent-4-ene, 5-(1,1-dimethylethyl)-2-methoxy-2-methyl-3-(2-phenylethenyl)-, $[2\alpha, 3\alpha(E)]$ (9CI) (CA INDEX NAME)

Relative stereochemistry.

Double bond geometry as shown.

- RN 174870-74-7 HCAPLUS
- CN 1-Oxa-2-silacyclohepta-4,6-diene, 7-(1,1-dimethylethyl)-2-methoxy-2-methyl-3-phenyl-, cis- (9CI) (CA INDEX NAME)

Relative stereochemistry.

- RN 174870-75-8 HCAPLUS
- CN 1-Oxa-2-silacyclohepta-4,6-diene, 7-(1,1-dimethylethyl)-2-methoxy-2-methyl-3-phenyl-, trans- (9CI) (CA INDEX NAME)

Relative stereochemistry.

174956-09-3 HCAPLUS

CN 1-0xa-2-silacyclopent-4-ene, 2-methoxy-2,4,5-trimethyl-3-(2phenylethenyl)-, [2α,3β(E)]- (9CI) (CA INDEX NAME)

Relative stereochemistry.

Double bond geometry as shown.

174956-11-7 HCAPLUS RN

1-Oxa-2-silacyclopent-4-ene, 5-(1,1-dimethylethyl)-2-methoxy-2methyl-3-(2-phenylethenyl)-, $[2\alpha, 3\beta(E)]$ - (9CI) (CA INDEX NAME)

Relative stereochemistry.

Double bond geometry as shown.

CC 29-6 (Organometallic and Organometalloidal Compounds) Section cross-reference(s): 22

174870-64-5P 174870-65-6P 174870-66-7P 174870-67-8P 174870-69-0P 174870-70-3P 174870-68-9P 174870-71-4P 174870-72-5P 174870-73-6P 174870-74-7P 174870-75-8P 174870-76-9P 174870-77-0P 174956-07-1P 174956-08-2P 174956-09-3P 174956-10-6P 174956-11-7P 174956-12-8P RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of)

L7 ANSWER 7 OF 11 HCAPLUS COPYRIGHT 2007 ACS on STN ACCESSION NUMBER: 1994:605453 HCAPLUS Full-text

DOCUMENT NUMBER: 121:205453

TITLE: Synthesis and reactivity of functional

1-methyl-silacyclopentenes AUTHOR(S): Heinicke, J.; Gehrhus, B.; Meinel, S.

CORPORATE SOURCE: Institut fuer Anorganische Chemie,

EMA-Universitaet Greifswald, Soldtmannstr. 16, Greifswald, D-17489, Germany

SOURCE: Journal of Organometallic Chemistry (1994), 474(1-2), 71-82

CODEN: JORCAI; ISSN: 0022-328X DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 121:205453

AB The silylenes MeSiCl, MeSiOMe and MeSiNMe2 can be thermally generated from disilanes and trapped by butadiene, isoprene and 2,3-dimethylbutadiene to give functionally-substituted silacyclopentenes. Some examples of cycloadins. to heterodienes are included to demonstrate the scope and mechanism of the reaction. The reactivity of 1-chloro-1-methyl-silacyclopent-3-ene has been studied.

IT 158172-38-4P

RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)

RN 158172-38-4 HCAPLUS

CN 1-Oxa-2-silacyclopent-4-ene, 2-methoxy-2,5-dimethyl-3-phenyl- (CA INDEX NAME)

```
29-6 (Organometallic and Organometalloidal Compounds)
ΙT
    100-40-3P, 4-Vinylcyclohexene 15983-92-3P 16054-12-9P
                 24429-79-6P 50694-31-0P 50694-33-2P 50694-36-5P 73357-64-9P 119820-51-8P 119820-52-9P
    24429-72-9P
    55544-25-7P
    158172-19-1P
                  158172-20-4P 158172-21-5P
                                               158172-22-6P
    158172-23-7P 158172-24-8P 158172-25-9P 158172-26-0P
    158172-27-1P 158172-28-2P 158172-29-3P 158172-30-6P
    158172-31-7P 158172-32-8P 158172-33-9P 158172-34-0P
    158172-35-1P 158172-36-2P 158172-37-3P 158172-38-4P
    158172-39-5P 158172-40-8P 158172-41-9P
                                                158172-42-0P
    158172-43-1P 158172-44-2P 158172-45-3P
                                                158172-46-4P
    RL: SPN (Synthetic preparation); PREP (Preparation)
```

L7 ANSWER 8 OF 11 HCAPLUS COPYRIGHT 2007 ACS on STN ACCESSION NUMBER: 1991:6591 HCAPLUS Full-text

DOCUMENT NUMBER: 114:6591

(preparation of)

TITLE: Silicon functionalized silyl enol ethers. 4.

Effects of variations of alkoxy substituents

upon cyclizations of dialkoxy-2-chloroethylsilyl

enol ethers to form 2,2-dialkoxy-1-oxa-2silacyclohexanes

AUTHOR(S): Walkup, Robert D.; Obevesekere, Nihal U.; Kane,

Robert R.

CORPORATE SOURCE: Dep. Chem. Biochem., Texas Tech. Univ., Lubbock,

TX, 79409-1061, USA

SOURCE: Chemistry Letters (1990), (7), 1055-8

CODEN: CMLTAG; ISSN: 0366-7022 DOCUMENT TYPE: Journal

LANGUAGE: English
OTHER SOURCE(S): CASREACT 114:6591

GT

- AB Various dialkoxychloroethylsilyl enol ether derivs. of pinacolone Me3CC(:CH2)OSi(OR)2CH2CH2Cl [R = Me, Et, (OR)2 = MeCHOCH2CHOMe, Me2COCH2CHOMe, Me2COCH2COMe2], prepared by lithiation of pinacolone with LiN(CHMe2)2 in Et20, silylation with Cl3SiCH2CH2Cl followed by alkoxylation, upon treatment with Bu3SnH and AIBN, underwent free-radical cyclizations to yield isolable 2,2dialkoxy-1-oxa-2- silacyclohexanes I. As the bulkiness of the alkoxy groups increased, the selectivity of the reaction for forming cyclized instead of directly reduced acyclic byproducts improved.
- 130891-57-5P 130891-58-6P

RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)

RN 130891-57-5 HCAPLUS

1-0xa-2-silacyclohexane, 6-(1,1-dimethylethyl)-2,2-dimethoxy- (CA INDEX NAME)

RN 130891-58-6 HCAPLUS

CN 1-0xa-2-silacyclohexane, 6-(1,1-dimethylethyl)-2,2-diethoxy- (CA INDEX NAME)

29-6 (Organometallic and Organometalloidal Compounds)

130891-57-5P 130891-58-6P 130891-59-7P

130891-60-0P 130891-61-1P 130891-63-3P 130891-64-4P 130891-65-5P 130891-66-6P 130891-67-7P 130891-68-8P

RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of) ANSWER 9 OF 11 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1981:405934 HCAPLUS Full-text

DOCUMENT NUMBER: 95:5934 TITLE: Silicon-containing heterocyclic compounds.

XXXVIII. Mass-spectrometric study of

1-oxa-2-silacvclohexa-3,5-diene derivatives AUTHOR(S): Bochkarev, V. N.; Polivanov, A. N.; Krasnova, T.

L.; Labartkava, M. O.; Silkina, N. N.;

Chernyshev, E. A.

CORPORATE SOURCE: Gos. Nauchno-Issled. Inst. Khim. Tekhnol. Elementoorg. Soedin., Moscow, USSR

SOURCE: Zhurnal Obshchei Khimii (1981), 51(1), 119-22

CODEN: ZOKHA4; ISSN: 0044-460X

DOCUMENT TYPE: Journal Russian

LANGUAGE:

AB The mass spectral fragmentation paths of I (R = H, Me, Ph; R1 = H, Me; R2 = Cl, Me, OEt) depended strongly on the substituents. Rearrangement processes were found and interpreted.

76470-24-1

RL: PRP (Properties) (mass spectrum of)

RN 76470-24-1 HCAPLUS

1-0xa-2-silacyclohexa-3,5-diene, 2,2-diethoxy-6-methyl- (9CI) (CA CN INDEX NAME)

22-2 (Physical Organic Chemistry)

IT 54905-28-1 67078-75-5 67608-54-2 69586-09-0 76470-24-1

77436-52-3 77436-53-4 RL: PRP (Properties)

(mass spectrum of)

L7 ANSWER 10 OF 11 HCAPLUS COPYRIGHT 2007 ACS on STN ACCESSION NUMBER: 1981:64674 HCAPLUS Full-text

DOCUMENT NUMBER: 94:64674

TITLE: Formation of pseudomolecular ions of compounds containing a silicon-carbon double bond under

electron impact

AUTHOR(S): Bochkarev, V. N.; Polivanov, A. N.; Slyusarenko,

T. F.; Bernadskii, A. A.

CORPORATE SOURCE: USSR

SOURCE: Zhurnal Obshchei Khimii (1980), 50(8), 1783-7

CODEN: ZOKHA4; ISSN: 0044-460X

DOCUMENT TYPE: Journal

Russian LANGUAGE:

- AΒ The mass spectra of several cyclic Si compds., e.g., I (R = Cl, Et, MeO, F) showed high-intensity peaks corresponding to species containing the Si:C group. Allylic and benzylic compds. also exhibited these peaks.
- 76470-24-1

RL: PRP (Properties) (mass spectrum of)

76470-24-1 HCAPLUS RN

CN 1-0xa-2-silacyclohexa-3,5-diene, 2,2-diethoxy-6-methyl- (9CI) (CA INDEX NAME)

```
22-2 (Physical Organic Chemistry)
ΙT
    107-37-9
              701-35-9
                         711-53-5
                                    770-10-5 872-46-8
                                                          999-94-0
    1112-55-6
               1745-72-8
                            2999-78-2
                                       4142-85-2 13093-11-3
    16054-12-9
                 18147-55-2
                              20151-87-5
                                           28871-17-2
                 38948-58-2
                              51986-73-3
                                           52023-18-4
    34106-93-9
                                                        54113-94-9
    54321-30-1
                 57831-44-4
                              61667-33-2
                                           67608-54-2
                                                       75722-33-7
    76470-24-1
                 76470-25-2
                              76470-26-3
    RL: PRP (Properties)
        (mass spectrum of)
```

ANSWER 11 OF 11 HCAPLUS COPYRIGHT 2007 ACS on STN ACCESSION NUMBER: 1976:180327 HCAPLUS Full-text DOCUMENT NUMBER: 84:180327

84:29235a,29238a ORIGINAL REFERENCE NO.:

TITLE: Copyrolysis of sym-tetramethoxydimethyldisilane

and 2,5-dimethvlfuran AUTHOR(S): Childs, Michael E.; Weber, William P.

CORPORATE SOURCE: Dep. Chem., Univ. South. California, Los

Angeles, CA, USA

SOURCE: Journal of Organic Chemistry (1976), 41(10),

1799-802

CODEN: JOCEAH: ISSN: 0022-3263

DOCUMENT TYPE: Journal

LANGUAGE: English

The reaction of methoxymethylsilylene (generated by pyrolysis of symtetramethoxydimethyldisilane) with 2,5-dimethylfuran yields 2-methoxy-2,3,6trimethyl-1-oxa-2-silacyclohexa-3,5-diene and 2-methoxy-2,4,7-trimethyl-1,3dioxa-2-silacyclohepta-4,6-diene.

58449-10-8P

RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)

RN 58449-10-8 HCAPLUS

CN 1-0xa-2-silacyclohexa-3,5-diene, 2-methoxy-2,3,6-trimethyl- (9CI) (CA INDEX NAME)

- CC 29-6 (Organometallic and Organometalloidal Compounds) Section cross-reference(s): 27, 28
- IT 58449-10-8P 58449-11-9P 58449-12-0P 58449-13-1P RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)

=>